

Recommended vaccinations for children up to 6 years old

If your child misses a shot, you don't need to start over, just go back to your child's doctor for the next shot. Talk with your child's doctor if you have any questions.

| Age | Vaccine & Dose | Age | Vaccine & Dose |
|----------------|--|--------------------|---|
| At birth | Hepatitis B (Dose 1 of 3) | 6 months or older | Influenza (Dose 1 of 2) Influenza (Dose 2 of 2 a month later and one dose yearly thereafter) |
| 1 to 2 months | Hepatitis B (Dose 2 of 3) | | |
| 2 months | DTaP (Dose 1 of 5) Hib (Dose 1 of 4) Polio (IPV) (Dose 1 of 4) Pneumococcal conjugate (PCV13) (Dose 1 of 4) Rotavirus (Dose 1 of 3) | 12 to 15 months | Hib (Dose 4 of 4) Pneumococcal conjugate (PCV13) (Dose 4 of 4) MMR (Dose 1 of 2) Varicella (Dose 1 of 2) |
| 4 months | DTaP (Dose 2 of 5) Hib (Dose 2 of 4) Polio (IPV) (Dose 2 of 4) Pneumococcal conjugate (PCV13) (Dose 2 of 4) Rotavirus (Dose 2 of 3) | 12 to 23 months | Hepatitis A (Dose 1 of 2) |
| 6 months | DTaP (Dose 3 of 5) Hib (Dose 3 of 4) Pneumococcal conjugate (PCV13) (Dose 3 of 4) Rotavirus (Dose 3 of 3) | 15 to 18 months | DTaP (Dose 4 of 5) |
| 6 to 18 months | Hepatitis B (Dose 3 of 3) Polio (IPV) (Dose 3 of 4) | 18 months or older | Hepatitis A (Dose 2 of 2, follows 6 months after Dose 1) |
| | | 4 - 6 years | DTaP (Dose 5 of 5) Polio (IPV) (Dose 4 of 4) MMR (Dose 2 of 2) Varicella (Dose 2 of 2) |

NOTE: Consult your child's pediatrician or health-care professional about catch-up vaccinations and vaccines for travelers or if your child has special health conditions.

SOURCE: U.S. Department of Health and Human Services